7.5 Subsequent Screening

Of the sites remaining after the WAG ERA HQ screening step all have nonradiological contamination and none have radiological contamination with HQ's greater than the target value.

Additional screenings and discussion are appropriate at this stage of the WAG ERA process to use a weight of evidence approach for the further elimination of sites and contaminants for consideration in the FS process. This type of evaluation will eliminate unnecessary and undesirable remediations for ecological receptors based on the following rationale.

- Contaminant concentrations—For the ERA, data values calculated for the BRA for the human health risk assessment were used when available. For those concentrations determined in the human health risk assessment (for the sites identified in Table 7-11) the maximum was generally used to estimate exposure-point concentrations except as noted.
- Modeling Conservatism—The exposure scenario used for ecological receptors assumes that the fences are down and the site has a viable habitat that is completely accessible to receptors. However, some sites of concern at CFA are currently fenced. Both the fence and the activities associated with this currently active facility should limit the exposure of receptors to much less than that model in the ERA. Additionally, (with some exceptions [particularly sites with water sources]) some of these sites are gravel and unsuitable habitat at the present time and would not provide any special attraction to ecological receptors.
- Parameter Conservatism—It is accepted in the risk assessment process that the many of the
 input parameters are developed to be conservatively protective of the receptors. Particularly,
 based on limited knowledge and the uncertainty of extrapolating to multiple species, TRVs
 development is very conservative. This is particularly true for naturally occurring metals,
 which can vary greatly across regions.

Based on this rationale, an additional screening was determined appropriate for the WAG 4 sites as agreed upon in a March 19, 1998 meeting between DOE-ID, EPA and IDHW. Contaminants are eliminated as a concern if the exposure point concentration is less than 10x the background value. For those contaminants that have no site-specific background the mean for the western states presented in Shacklette and Boerngen (1984) or other sources was considered acceptable. Results of this screening are presented in Table 7-23.

This screening resulted in eliminating one site (CFA-06). The remaining 13 sites (CFA-01, CFA-02, CFA-04, CFA-05 CFA-08, CFA-10, CFA-13, CFA-21, CFA-26, CFA-40, CFA-41, CFA-43, and CFA-51) will be evaluated further in the FS.

7.6 Transition to Sitewide ERA

This WAG ERA represents the second phase of the four-phased approach to ERA presented in Section 7.1 (see Figure 7-1). The approach applies an iterative, "tiered" process in which preliminary assessments, based on conservative assumptions, support progressively more refined assessments (Maughn 1993; Opresko et al. 1995; Levin et al. 1989).

The first phase includes a data review and either a SLERA or EDGA, which is a "pre-assessment" performed at the WAG level. The pre-assessment (1) reduces the number of contaminants and sites to be

 Table 7-23. Results of WAG 4 ecological contaminant screening against 10x INEEL background concentrations.

Site Number	Site Description and Size (sq. meters)	Contaminant of Potential Concern	Hazard Quotient	Maximum Concentration (mg/kg)	10x INEEL Background* (mg/kg)	Comment	Retain COPC?
CFA-01	Landfill I	Benzo(a)pyrene	<1 to 2	0.89	NA	No background	YES
	4.30E+04	Chrysene	<1 to 20	450	NA	No background	YES
		Chromium III	<1 to 50	53	330	<10x background	No
		Copper	<1 to 30	73.4	220	<10x background	No
		Lead	≤1 to 200	97	170	<10x background	No
			<1 to 6				
		Mercury		0.08	0.5	< 10x background	No
		Silver	≤1 to 10	19.5	NA	No background	YES
		Zinc	≤1 to 30	230	1500	<10x background	No
CFA-02	Landfill II 7.07E+05	4-methyl-2-pentanone	NA	0.02	NA	No background	YES
		2-methylnapthalene	NA	0.05	NA	No background	YES
		Acetone	≤1 to 20	5.8	NA	No background	YES
		Arsenic	≤1 to 20	17	58	<10x background	No
		Dibenzofuran	NA	0.039	NA	No background	YES
		Lead	950	255	170	>10x background	YES
		Mercury	≤1 to 20	0.19	0.5	<10x background	No
			_1 10 20	<i>-</i>	-	va outle- outle	No
		Pentachlorophenol	NA	0.074	NA	No background	YES

-12

Table 7-23. (continued).

Site Number	Site Description and Size (sq. meters)	Contaminant of Potential Concern	Hazard Quotient	Maximum Concentration (mg/kg)	10x INEEL Background* (mg/kg)	Comment	Retain COPC?
CFA-04	Pond near CFA-674	Arsenic	<1 to 20	22.4	58	<10x background	No
	6.88E+03	Barium	≤1 to 6	1120	3,000	<10x background	
		Cadmium	≤1 to 1,000	3.4	22	<10x background	No
		Chromium III	<1 to 100	237	330	<10x background	No
		Cobalt	≤1 to 20	10	110	<10x background	No
		Copper	≤1 to 150	365	220	<10x background	YES
		Lead	≤1 to 90	42.4	170	<10x background	No
		Mercury	<1 to 30,000	439	0.5	>10x background	YES
		Nickel	<1 to 110	355	350	<10x background	YES
		Silver	<1 to 20	31	NA	No background	YES
		Vanadium	≤2 to 200	46	450	<10x background	No
CFA-05	Motor Pool Pond 7.43E+03	4-methyl-2-pentanone	NA	0.065	NA	No background	YES
		Arsenic	≤1 to 5	19.8	58	<10x background	No
		Cadmium	≤1 to 4,000	38.8	22	>10x background	YES
		Chromium III	≤1 to 90	91.3	330	<10x background	No
		Cobalt	≤2 to 20	15.0	110	<10x background	No
		Copper	≤1 to 40	342	220	>10x background	YES
		Lead	≤1 to 200	464	170	>10x background	YES
		Manganese	≤1 to 14	617	4,900	<10x background	No
		Mercury	≤1 to 9	0.58	0.5	>10x background	YES

7-123

Table 7-23. (continued).

Site Number	Site Description and Size (sq. meters)	Contaminant of Potential Concern	Hazard Quotient	Maximum Concentration (mg/kg)	10x INEEL Background* (mg/kg)	Comment	Retain COPC?
		Nickel	≤1 to 3	36.7	350	<10x background	No
		Selenium	<1 to 5	1.1	2.2	<10x background	No
		Vanadium	<1 to 20	34	450	<10x background	No
		Zinc	<1 to 20	241	1500	<10x background	No
CFA-06	Lead Shop (outside areas) 2.50E+03	Arsenic	≤1 to 10	14.5	58	<10x background	No
		Lead	≤1 to 200	153	170	<10x background	No
CFA-08	Sewage Plant (CFA-691), Septic Tank (CFA-716), and Drainfield 1.85E+04	Arsenic	≤1 to 10	16.8	58	<10x background	No
		Chloromethane	NA	0.005	NA	No background	YES
		Chromium III	<1 to 40	77.6	330	<10x background	No
		Соррег	≤1 to 20	56.5	220	<10x background	No
		Lead	≤1 to 40	18	170	<10x background	No
		Mercury	≤1 to 30	0.51	0.5	>10x background	YES
		Nickel	≤1 to 20	90.3	350	<10x background	No
		Selenium	≤1 to 7	0.5	2.2	<10x background	No
		Silver	≤1 to 10	24.1	NA	No background	YES
CFA-10	Transformer Yard Oil Spills 8.08E+02	Antimony	<1 to 4	9.5	48	<10x background	No
		Arsenic	<1 to 8	11.6	58	<10x background	No

7 - 122

Table 7-23. (continued).

Site Number	Site Description and Size (sq. meters)	Contaminant of Potential Concern	Hazard Quotient	Maximum Concentration (mg/kg)	10x INEEL Background* (mg/kg)	Comment	Retain COPC?
		Cadmium	≤1 to 2,000	7.3	22	<10x background	No
		Chromium III	<1 to 100	102	330	<10x background	No
		Cobalt	≤1 to 30	15.7	110	<10x background	No
		Copper	<1 to 70	259	220	>10x background	YES
		Lead	<1 to 5000	5,560	170	>10x background	YES
		Manganese	≤ 1 to 20	509	4,900	<10x background	No
		Mercury	<1 to 4	0.09	0.5	<10x background	No
		Nickel	<1 to 20	111	350	<10x background	No
		Zinc	<1 to 70	1,150	1,500	<10x background	No
CFA-13	Dry Well (South of	Antimony	<1 to 3	11.5	48	<10x background	No
	CFA-640) 2.50E+01	Cadmium	<1 to 60	7.37	22	<10x background	No
		Chromium III	<1 to 300	267	330	<10x background	No
		Copper	≤1 to 20	1,900	220	>10 background	YES
		Lead	<1 to 20	725	170	>10 background	YES
		Mercury	<1 to 7	1.97	0.5	>10 background	YES
		Nickel	<1 to 3	85.1	350	<10x background	No
		Silver	<1 to 10	19.4	NA	No background	YES
		Zinc	<1 to 6	302	1500	<10x background	No

/-12:

Table 7-23. (continued).

Site Number	Site Description and Size (sq. meters)	Contaminant of Potential Concern	Hazard Quotient	Maximum Concentration (mg/kg)	10x INEEL Background* (mg/kg)	Comment	Retain COPC?
CFA-21	Fuel Tank at Nevada Circle (S by CFA-629) 7.00E+00	ТРН	<1 to 3	54,000	NA	No background	YES
CFA-26	CFA-760 Pump Station Fuel Spills 1.12E+02	ТРН	≤1 to ≤4	3,470	NA	No background	YES
CFA-40	Returnable Drum Storage (south of CFA-601) 5.40E+02	ТРН	<1 to 3	<625	NA	No background	YES
CFA-41	Excess Drum Storage (south of CFA-674) 6.97E+03	ТРН	<1 to 20	<1,000	NA	No background	YES
CFA-43	Lead Storage Area 1.53E+04	Lead	≤1 to 300	180	170	>10x background	YES
CFA-51	Dry Well at north end of CFA-640 1.00E-01	Cadmium	<1 to 5	14.0	22	<10x background	No
		Copper	<1 to3	250	220		YES

/-120

Table 7-23. (continued).

Site Number	Site Description and Size (sq. meters)	Contaminant of Potential Concern	Hazard Quotient	Maximum Concentration (mg/kg)	10x INEEL Background* (mg/kg)	Comment	Retain COPC?
		Zinc	<1 to 7	340	1500	<10x background	No
At CFA-17/47	organic compounds were ca	alculated using maximum concen	itrations.				
Xylene was det	ected at 10 ft bgs.						
Metal concentra	ations (mg/kg) typically fou	and in INEEL soil is discussed in	Rood et al. 1996				

addressed in the WAG-level ERA by eliminating those that clearly pose a low likelihood for risk, (2) better defines the nature and extent of contamination at individual WAG sites, (3) indicates sites for which further data are needed, and (4) identifies other data gaps. The results of this assessment serve to support problem formulation and drive media and pathways to be evaluated for the WAG-level ERA. This pre-assessment methodology will be used to screen additional sites and/or contaminants identified during the ongoing CERCLA process. This level of the assessment does not support setting remedial action levels. Details of SLERA methodology can be found in VanHorn et al. (1995).

The second phase is the WAG ERA, which represents the baseline risk assessment of the No Action alternative for remediation. The WAG ERA incorporates the screening results to assess potential risks to ecological receptors at the WAG-level following EPA guidance (1992). The method parallels the human health risk assessment in that each site at the WAG is individually assessed. This section presents the WAG ERA for OU 4-13. Table 7-21 summarizes the ERA screening steps used at WAG 4 sites. The WAG ERA results will (1) provide a list of COPCs to be addressed in the OU 10-04 ERA, and, (2) identify OU 10-04 level data gaps that must be filled before performing the OU 10-04 ERA. The results of the WAG ERA may also support risk assessments to evaluate WAG remedial actions or additional assessments, if necessary.

The third phase of the ERA process is the OU 10-04 ERA, which is performed to evaluate risk to INEEL-wide ecological resources. The OU 10-04 ERA will integrate the results of the WAG ERAs for all INEEL WAGs to determine whether contamination at the WAGs contributes potential risk to populations and communities on an ecosystem-wide basis (i.e., over the entire INEEL).

7.7 References

- Abbey, H. K. and N. Platonow, 1968, "Toxicity of Vanadium in Calves". Vet. Record, 82:292.
- Adams, W. J., R. A. Kimberle, and J. W. Barnett, Jr., 1992, "A Sediment Quality and Aquatic Life Assessment," *Environmental Science and Technology*, 26(10):1865–1875.
- Adriano, D. C., 1986, Trace Elements in the Terrestrial Environment, Springer-Verlag, New York, NY.
- Agency for Toxic Substances and Disease Registry (ATSDR), 1993. Draft: Toxicological Profile for Cadmium, April.
- Albanus, L., L., et al, 1972, "Toxicity for Cats of Methylmercury in Contaminated Fish From Swedish Lakes and of Methylmercury Hydroxide Added to Fish," *Environmental Research*, 5(4):425-42.
- Ambrose, A. M., et al., 1976, "Long-term Toxicologic Assessment of Nickel in Rats and Dogs," *Journal of Food Science and Technology*, 13: 181-187.
- American Biogenics Corp., 1986, Ninety-day Gavage Study in Albino Rats Using Nickel, Draft Final Report submitted to Research Triangle Institute, P.O. Box 12194, Research Triangle Park, NC 27709.
- Ammerman, C. B. et al. 1973, Toxicity of Certain Minerals to Domestic Animals. A Review. Fla. Agric. Exp. Stn. Res. Bull. AL73-6. University of Florida, Gainsville.
- Anwar, R. A., et al., 1961, "Chronic toxicity studies. III, Chronic Toxicity of Cadmium and Chromium in Dogs," Archive of Environmental Health, 3:456-467.

- Arthur D. Little, Inc. (ADL), 1987, *The Installation Restoration Program Toxicology Guide*, Vols. I-III, Cambridge, MA.
- Arthur, W. J. and O. D. Markham, 1983, "Small Mammal Soil Burrowing as a Radionuclide Transport Vector at a Radioactive Waste Disposal Area in Southeastern Idaho," *Health Physics*, 43:493–500.
- Arthur, W. J. et al., 1984, Vertebrates of the Idaho National Engineering Laboratory, DOE/ID-12099, U.S. Department of Energy Idaho Operations Office, Radiological and Environmental Sciences Laboratory.
- Arthur, W. J., 1987, "Radiation Dose to Small Mammals Inhabiting a Solid Radioactive Waste Disposal Area in Southeastern Idaho," *Journal of Applied Ecology*, 23:13–26.
- Arthur, W. J., and R. J. Gates, 1988, "Trace Element Intake via Soil Ingestion in Pronghorns and in Black-tailed Jack Rabbits," *Journal of Range Management*, 41:162–166.
- ASTDR (Agency for Toxic Substances and Disease Registry), 1990b, Toxicological Profile for Benzo(b)fluoranthene, March.
- ATSDR, 1988a, Toxicological Profile for Nickel, PB89-160378, December.
- ATSDR, 1990a, Draft Toxicological Profile for Cobalt, October.
- ATSDR, 1990c, Toxicological Profile for Chrysene, March.
- ATSDR, 1991, Draft Toxicological Profile for Boron, February 15.
- ATSDR, 1992, Agency for Toxic Substances and Disease Registry, *Toxicological Profile for Acetone*, February.
- ATSDR, 1992, Toxicological Profile for Barium and Compounds, PB93-110658, July.
- ATSDR, 1992, *Toxicological Profile for Vanadium*, Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.
- Aulerich, R. J., R. K. Ringer, and J. Iwamoto, 1974, "Effects of Dietary Mercury on Mink," Archives of Environmental Contamination and Toxicology, 2:43–51.
- Baes, C. F. and Sharp, R. D., 1983, "Proposal for Estimation of Soil Leaching and Leaching Constants for Use in Assessment Models," *J. Environmental Quality*, 12(1):17–28.
- Baes, C. F., III, et al., 1984, A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides Through Agriculture, ORNL-5786, U.S. Department of Energy. Oak Ridge National Laboratory, Oak Ridge, TN.
- Barnthouse, L. W., G. W. I. Suter, and A. E. Rosen, 1990, "Risk of Toxic Contaminants to Exploited Fish Populations: Influence of Life History, Data Uncertainty and Exploitation Intensity," *Environmental Toxicology and Chemistry*, 9:297–311.

- Barnthouse, L. W., Suter, G. W., II, Bartell, S. M., Beauchamp, J. J., Gardner, R. H., Linder, E., O'Neill, R. V., and Rosen, A. E. (1986) User's Manual for Ecological Risk Assessment. Environmental Sciences Division Publication No. 2679, Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Bartlett, R. J. and J. M. Kimble, 1976, "Behavior of Chromium in Soils: II. Hexavalent Forms," J. Environ. Qual., 5(1):383-386.
- Bartlett, R. J., 1991, "Chromium Cycling in Soils and Water: Links, Gaps, and Methods," *Environmental Health Perspectives*, 92:17–24.
- Beyer, W. N., E. E. Conner, and S. Geroud, 1994. "Estimates of Soil Ingestion by Wildlife," *Journal of Wildlife Management*, 58, pp. 375–382.
- Bird, D. M., et al., 1983, "Synergistic Effects of Aroclor 1254 and Mirex on the Semen Characteristics of American Kestrels," Archives of Environmental Contamination and Toxicology, 12:633-640.
- Borg, K., K. et al, 1970, "Experimental Secondary Methyl Mercury Poisoning in the Goshawk, Accipiter G. gentilis L.," Environmental Pollution, 1:91-104.
- Bosch, H. M., A. B. Rosefield, R. Houston, H. R. Shipman, and F. L. Woodward, 1950, Methemoglobinemia and Minnesota Well Supplies, J. Am. Water Works Association, 42:161-170.
- Brewer, B., 1940, "A statistical study of cobalt polycythemia in the dog," Am. J. Physiol. 128:345–348.
- Brookins, D. G., 1988, Eh-pH Diagrams for Geoechemistry, Springer-Verlag, New York, NHY, p. 373.
- Brune et al., 1981, "Investigation of the Tumorigenic Response of Benzo(a)pyrene in Aqueous Caffeine Solution Applied Orally to Sprague-Dawley Rats," *Journal of Cancer Research and Clinical Oncology*, 102(2):153–157.
- Burt, W. H., and R. P. Gotssenheider, 1976, A Field Guide to the Mammals of America North of Mexico, Houghton Mifflin, Boston.
- Byerrum, R. U. et al. 1974. Vanadium. Washington, D.C.: National Academy of Sciences, 1-117.
- Byron, W. R., G. W. Bierbower, J. B. Brouwer, and W. H. Hansen, 1967, "Pathologic Changes in Rats and Dogs from Two-year Feeding of Sodium Arsenite or Sodium Arsenate," *Toxicology and Applied Pharmacology*, 10:132–147.
- Cakir, A., T. W. Sullivan, and F. B. Mather, 1978, Alleviation of Fluorine Toxicity in Starting Turkeys and Chicks with Aluminum, *Poultry Science*, 57:498, as cited in National Academy of Sciences, 1980, Mineral Tolerance of Domestic Animals, Washington, D.C.
- Callahan, M. A. et al., 1979, Water-Related Environmental Fate of 129 Priority Pollutants, EPA 440/4-79-029a, Office of Water Planning and Standards, U.S. Environmental Protection Agency, Washington, D.C.

- Carey, E. E., 1982, "Chromium in Air, Soil and Natural Waters," *Topics in Environmental Health 5, Biological and Environmental Aspects of Chromium, S. Langar (ed.), Elsevier Publishers, New York, NY, pp. 49-64.*
- CDC, 1994, Rare, Threatened, and Endangered Plants and Animals of Idaho, Third edition, Idaho Department of Fish and Game, Boise, Idaho, 39p.
- Chang, C. W. J., R. M. Nakamura, and C. C. Brooks, 1977, "Effect of Varied Dietary Levels and Forms of Mercury on Swine," J. Anim. Sci., 45:279.
- Charbonneau, S. M. et al., 1976, "Chronic Toxicity of Methylmercury in the Adult Cat, Interim Report," *Toxicology*, 5(3):337-49.
- Cholewa, A. F. and LD. M. Henderson, 1984, A Survey and Assessment of Rare Vascular Plants of the Idaho National Engineering Laboratory Site, DOE/ID-12100, Radiological and Environmental Sciences Laboratory, Idaho Falls, ID, pp. 45.
- Cieminski, K. L., 1993, Wildlife Use of Wastewater Ponds at the Idaho National Engineering Laboratory, M.S. Thesis, South Dakota University, Brookings, 233p.
- Cohen, M. D. et al., 1993, "Mechanisms of Chromium Carcinogenicity and Toxicity," *Critical Review of Toxicology*, 23(3):255–281.
- Colle, A., et al. 1980, "Lead Poisoning in Monkeys: Functional and Histopathological Alterations of the Kidneys," *Toxicology*, 18:145–158.
- Coughtrey, P. J., Jackson, D., and Thorne, M. C., 1985, Radionuclide Distribution and Transport in Terrestrial and Aquatic Ecosystems: A Compendium of Data, Netherlands, A.A. Balkema.
- Dahlgren, R. B., and R. L. Linder, 1971, "Effects of Polychlorinated Biphenyls on Pheasant Reproduction, Behavior, and Survival," *Journal of Wildlife Management*, 35(2):315–319.
- Dahlgren, R. B., R. L. Linder, and C. W. Carlson, 1972, "Polychlorinated Biphenyls: Their Effects on Penned Pheasants," *Environmental Health Perspectives*, 1:89–101.
- Del Debbio, J. A. and Thomas, T. R., 1989, Transport Properties of Radionuclides and Hazardous Chemical Species in Soils at the Idaho Chemical Processing Plant, WINCO-1068.
- DeMayo, A., M. et al., 1982, "Toxic Effects of Lead and Lead Compounds on Human Health, Aquatic Life, Wildlife, Plants, and Livestock," CRC Critical Review Environmental Control, 12:257–305.
- Dieter, M. P and M. T. Finley, 1978, "Erythrocyte Gamma-aminolevulinic Acid Dehyratase Activity in Mallard Ducks: Duration of Inhibition After Lead Shot Dosage," *Journal of Wildlife Management*, 42:621–625. As cited in Eisler, 1988.
- Diller, L. V. and D. R. Johnson, 1988, Food Habits, Consumption Rates, and Predation Rate of Western Rattlesnakes and Gopher Snakes in Southwestern Idaho, Herpetologica 44(2):228-233.

- DOE-ID. Department of Energy, Idaho Field Office, 1991, Federal Facility Agreement and Consent Order for the Idaho National Engineering Laboratory, State of Idaho Department of Health and Welfare, U.S. Environmental Protection Agency, U.S. Department of Energy, December 4.
- DOE, 1993, INEL Stormwater Pollution Prevention Plan for Industrial Activities Storm Water General Permit #IDR00A194, DOE-ID-10431, Revision 3.
- DOE, 1994, Incorporating Ecological Risk Assessment into Remedial Investigation/Feasibility Study Work Plans, U.S. Department of Energy Memorandum, July.
- DOE-ID, 1993, Track 2 Sites: Guidance for Assessing Low Probability Hazard Sites at the INEL (draft), DOE/ID-10389, August.
- Donaldson, R. M. and R. F. Barreras, 1966, "Intestinal Absorption of Trace Quantities of Chromium," Journal of Laboratory and Clinical Medicine, 68:484-493.
- Dragun, J., 1988, The Soil Chemistry of Hazardous Material, Hazardous Materials Control Research Institute.
- Duce, R. A. and G. L. Hoffman. 1976. Atmospheric vanadium transport to the ocean. Atmospheric Environment 10:989–996.
- Dunning, J. B., Jr., 1993, CRC Handbook of Avian Body Masses, CRC Press, Boca Raton, FL.
- Eastin, W. C., Jr. and T. J. O'Shea, 1981, "Effects of Dietary Nickel on Mallards," *Journal of Toxicology and Environmental Health*, 7(6):883-892.
- Eaton, R. D., D. C. Secord, and P. Hewitt, 1980, "An Experimental Assessment of the Toxic Potential of Mercury in Ringed Seal Liver for Adult Laboratory Cats," *Toxicology and Applied Pharmacology*, 55(3):514-21.
- EG&G Idaho, Inc., 1993, Data Characterization Process in Environmental Restoration, Program Directive 3.7, EG&G Idaho, Idaho Falls, Idaho.
- Eisenbud, M., 1987, Environmental Radioactivity: From Natural, Industrial, and Military Sources, Third edition, Academic Press, New York, NY, 475 pp.
- Eisler, R., 1985a, Cadmium Hazard to Fish, Wildlife, and Invertebrates: A Synoptic Review, U.S. Fish and Wildlife Service, Biological Report, 85(1.2)
- Eisler, R., 1985b, Selenium Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, Fish and Wildlife Service, Biological Report No. 85 (1.5).
- Eisler, R., 1986a, Chromium Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, U.S. Fish and Wildlife Service Biological Report, 85(1.6).
- Eisler, R., 1986b, Polychlorinated biphenyls Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, U.S. Fish and Wildlife Service Biological Report, 7, April.

- Eisler, R., 1987, Mercury Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, U.S. Fish and Wildlife Service Biological Report, 85(1.10).
- Eisler, R., 1988a, Arsenic Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, U.S. Fish Wildlife Service Biological Report, 85(1.12): 92 pp.
- Eisler, R., 1988b, Lead Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, Fish and Wildlife Service Biological Report No. 85 (1.14).
- EPA, 1976, Quality Criteria for Water, EPA-440/9-76/023, U. S. Environmental Protection Agency, Washington, D.C.
- EPA, 1984, *Health Assessment for Toluene*, EPA-540/1-86-033. Office of Health and Environmental Assessment, Cincinnati, Ohio.
- EPA, 1985, Environmental Profiles and Hazard Indices for Constituents of Municipal Sludge: Office of Water Regulation and Standards, June.
- EPA, 1986b, Ninety-Day Gavage Study in Albino Rats Using Acetone. Office of Solid Waste, Washington, D.C.
- EPA, 1988, Guidance for Conducting Remedial Investigations and Feasibility Studies Under CERCLA, Office of Emergency and Remedial Response, Washington, D.C., (EPA540 G-89004), October.
- EPA, 1988a, Joint Memorandum of the Office of Emergency and Remedial Response and the Office of Waste Programs Enforcement, (cited in EPA, 1991).
- EPA, 1988b, Review of Ecological Risk Assessment Methods, EPA/230/10-88-041. Office of Planning and Evaluation, Washington, D.C.
- EPA, 1990, Transuranium Elements, Volume 1: Elements of Radiation Protection, EPA/520/1-90-015, Washington, D.C.
- EPA, 1992a, Framework for Ecological Risk Assessment, EPA/630/R-92/001, PB93-102192, U.S. Environmental Protection Agency, ORD/Risk Assessment Forum, February, 55 pp.
- EPA, 1992b, Supplemental Guidance to RAGS: Calculating the Concentration Term, Office of Solid Waste and Emergency Response, Washington, D.C., EPA Publication 9285.7-081, May.
- EPA, 1993, Great Lakes water quality initiative criteria documents for the protection of wildlife (proposed): DDT, Mercury, 2,3,7,8-TCDD, PCBs, EPA/822/R-93-007, U.S. Environmental Protection Agency, Office of Science and Technology, Washington, D.C.
- EPA, 1993a, Wildlife Exposure Factors Handbook, U.S. Environmental Protection Agency, Office of Research and Development, Washington, D.C., EPA/600/R-93/187B, December.
- EPA, 1993b, Ch. 9. Selenium Effects at Kesterson Reservoir, A Review of Ecological Assessment Case Studies from a Risk Assessment Perspective, EPA/630/R-92/005.

- EPA, 1997, Ecological Risk Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessment (Interim Final), U.S. Environmental Protection Agency, Environmental Response Team, Edison, NJ, June.
- Faure, G., 1991, *Principles and Applications of Inorganic Geochemistry*, MacMillan Publishing Company, New York, NY, 626.
- Fimreite, N., 1979, "Accumulation and Effects of Mercury on Birds," In *The Biogeochemistry of Mercury in the Environment*, J. O. Nriagu, Ed., Elsevier/North Holland Biomedical Press, New York.
- Fiussello, N. and P. Novo, 1977, "The Effect of Chromium on Plants," Allionia, 21:105-114.
- Franchini, I. and A. Mutti, 1988, "Selected Toxicological Aspects of Chromium(VI) Compounds," *The Science of the Total Environment*, 71(3):379-87.
- Friedlander, G. et al., 1981, Nuclear and Radiochemistry, 3rd Edition, John Wiley & Sons, NY, NY.
- Frost, D. V., 1983, "What Do Losses in Selenium and Arsenic Bioavailability Signify for Health?" *The Science of the Total Environment*, 28:455–466.
- Gabler, K. I., 1997, Distribution and Habitat Requirements of the Pygmy Rabbit (*Bracheylagus idahoensis*) on the Idaho National Engineering and Environmental Laboratory, M.S. Thesis (unpublished), Idaho State University.
- Ganther, H. E. 1980, "Interactions of Vitamin E and Selenium with Mercury and Silver," Annals of the New York Academy of Science, 355: 212-26.
- Gerber, G. B., J. Maes, and B. Eykens, 1982, "Transfer of Antimony and Arsenic to the Developing Organism," *Archives of Toxicology*, 49:159–168.
- Griffin, S. and P. Turck, 1991 "Bioavailability in Rabbits of Sodium Arsenite Absorbed to Soils," *The Toxicologist*, 11:718.
- Halverson, A. W., I. S. Palmer, and P. L. Guss. 1966, "Toxicity of Selenium to Post-weanling Rats," *Toxicol. Appl. Pharmacol.*, 9:477–484.
- Hampton, N. L. et al., 1995, A Preliminary Survey of the National Wetlands Inventory as Mapped for the Idaho National Engineering Laboratory, INEL-95/0101, February.
- Hawkins, D. B. and Short, H. L., 1965, Equations for the Sorption of Cesium and Strontium on Soil and Clinoptilolite, IDO-12046.
- Hecht, S. S., Bondinell, W. E., Hoffmann, D., 1974. "Chrysene and Methylchrysenes: Presence in Tobacco Smoke and Carcinogenicity" *J. Nat. Cancer Inst.*, 53:1121–1133.
- Heinz, G. H. and Haseltine, S. D., 1983, "Altered Avoidance Behavior of Young Black Ducks Fed Cadmium," *Environmental Toxicology and Chemistry*, 2:419-421. As cited in Eisler, 1985.
- Heinz, G. H. et al., 1987, "Reproduction of Mallards Fed Selenium," *Environmental Toxicology and Chemistry*, 6:423–433.

- Herigstad, R. R. et al., 1972, "Chronic Methylmercury Toxicosis in Calves," J. Am. Vet. Med. Associ., 160:173.
- Hill, C. H., 1979, "The effect of dietary protein levels on mineral toxicity in chicks," *Journal of Nutrition*, 109:501–507.
- Hill, C. H., 1979, "The Effect of Dietary Protein on Mineral Toxicity in Chicks," *Journal of Nutrition*, 109:501-507.
- Hilliard, H. E. 1987. Vanadium. The minerals yearbook minerals and metals. 917-927.
- Hoffman, D. J. et al., 1985, "Survival, Growth, and Accumulation of Ingested Lead in Nestling American Kestrels (Falco sparverius)," Comparative Biochemistry and Physiology, 80C: 431–439.
- Hoover, R. L., and D. L., Wills, eds., 1987, *Managing Forest Lands for Wildlife*, Developed in cooperation with the U.S. Department of Agriculture, Forest Service, Rocky Mountain Region. Colorado Division of Wildlife.
- Hopkin, S. P. and C. A. C. Hames, 1994, "Zinc, Among a 'Cocktail' of Metal Pollutants, is Responsible for the Absence of the Terrestrial Isopod *Porcellio scaber* from the Vicinity of a Primary Smelting Works," *Ecotoxicology*, 2:68–78.
- Hornshaw, et al., 1986, "LC50 Test Results in Polychlorinated Biphenyl-fed Mink: Age, Season, and Diet Comparisons," Archives of Environmental Contamination and Toxicology, 15(6):717-23.
- Hornshaw, T. C., R. J. Aulerich, and H. E. Johnson, 1983, "Feeding Great Lakes Fish to Mink: Effect on Mink and Accumulation and Elimination of PCBs by Mink," *Journal of Toxicology and Environmental Health*, 11:933–946.
- Huck, D. W. and A. J. Clawson, 1976, "Excess Dietary Cobalt in Pigs," J. Anim. Sci. 43:1231.
- Hugot, D., J. Causeret, and C. Richir, 1980, "The Influence of Large Amounts of Sodium Nitrate on the Reproductive Performances in Female Rats," Ann. Nutr. Alim. 34:1115–1124.
- IAEA (International Atomic Energy Agency), 1992, Effects of Ionizing Radiation on Plants and Animals at Levels Implied by Current Radiation Standards, Technical Report Series No. 332.
- IDFG (Idaho Department of Fish and Game), 1994, "Rare, Threatened and Endangered Plants and Animals of Idaho," Conservation Data Center, Boise, ID.
- INPS (Idaho Native Plant Society), 1995, Results of the Eleventh Annual Idaho Rare Plant Conference, February 7--8, Boise, ID.
- INPS (Idaho Native Plant Society), 1996, Results of the Twelfth Annual Idaho Rare Plant Conference, February 13–14, Boise, ID.
- Ivankovic, S. and R. Preussmann, 1975, "Absence of Toxic and Carcinogenic Effects after Administration of High Doses of Chromic Oxide Pigment in Subacute and Long-term Feeding Experiments in Rats," Food and Cosmetic Toxicology, 13(3):347–352.

- Kanisawa, M. and H. A. Schroeder, 1969, "Life Term Studies on the Effect of Trace Elements of Spontaneous Tumor in Mice and Rats," *Cancer Research*, 29(4):892–895.
- Keck, K. N., I. Porro, A. J. Sondrup, S. H. McCormick, S. M. Lewis, 1994, Remedial Investigation/Feasibility Study Report for Operable Unit 4-12: Central Facilities Area Landfills I, II and III at the Idaho National Engineering Laboratory, INEL-94/0124, November.
- Keller, B. L., W. R. Bosworth, and R. W. Doering, 1993, *Bat Habitat Research; Final Technical Report*, U.S. Department of Energy, Idaho Field Office, Idaho Falls. 20 pp.
- Khera, K. S. and S. A. Tobacova, 1973, "Effects of Methylmercuric Chloride on the Progeny of Mice and Rats Treated Before or During Gestation," *Food and Cosmetic Toxicology*, II:245-254.
- Khera, K. S., 1973, "Reproductive Capability of Male Rats and Mice Treated with Methylmercury," *Toxicology and Applied Pharmacology*, 24:167.
- Khera, K. S., 1979, Teratogenic and Genetic Effects of Mercury Toxicity, The Biogeochemistry of Mercury in the Environment, J. O. Nriagu, Ed., Elsevier/North-Holland Biomedical Press, New York, NY
- Kimmel, C. A. et al., 1980, "Chronic Low Level Lead Toxicity in the Rat. I. Maternal Toxicity and Perinatal Effects," *Toxicology and Applied Pharmacology*, 56:28–41.
- King, L. D., 1988, "Retention of Metals by Several Soils of the Southeastern United States," *J. Environ. Qual.* 17:239-246.
- Kirchman, R. et al., 1993, "Terrestrial Pathways," *Radioecology After Chernobyl*, E. E. Warner and R. M. Hamson (eds.), John Wiley and Sons, New York, NY.
- Klein, M., 1963, "Susceptibility of Strain B6AF/j Hybrid Infant Mice to Tumorigenesis with 1,1-benzoanthracene, Deoxycyclic Acid, and 3-methylcholanthrene," *Cancer Research*, 23:1701-1707.
- Kocher, 1981, Radioactive Decay Data Tables, NTIS DOE-TIC-11026.
- Kopp, J. F., 1969, The Occurrence of Trace Elements in Water, In: Hemphill D. D., ed., Trace Substances in Environmental Health, III. Proceedings of the University of Missouri Third Annual Conference on Trace Substances in Environmental Health. Columbia, MO: University of Missouri, 59.
- Kopp, S. J., et al., 1985, "Cardiovascular Dysfunction and Hypersensitivity to Sodium Pentobarbital Induced by Chronic Barium Chloride Ingestion," *Toxicology and Applied Pharmacology*, 77(23):303-314.
- Koval, P. J., T. J. Peterle, and J. D. Harder, 1987, "Effects of Polychlorinated Biphenyls on Mourning Dove Reproduction and Circulating Progesterone Levels," *Bulletin of Environmental Contamination and Toxicology*, 39:663-670.

- Kramber, W. J. et al., 1992, "Producing a Vegetation Map of the Idaho National Engineering Laboratory Using Landsat Thematic Mapper Data," *Proceedings of ASPRS 1992 Annual Meeting*, Albuquerque, NM, March.
- Kubena, L. F. and T. D. Philips, 1982, "Toxicity of vanadium in female leghorn chickens," Poultry Science, 62:47–50.
- Lane, R. W., B. L. Riddle, and J. F. Borzelleca, 1982, "Effects of 1,2 dichloroethane and 1,1,1-trichloroethane in Drinking Water on Reproduction and Development in Mice," *Toxicol. Appl Pharmacol.* 63:409–421.
- Levin, S. A., et al., 1989, *Ecotoxicology: Problems and Approaches*, Springer-Verlag, New York, New York, 547 pp.
- Lewis, S. C., J. R. Lynch, and A. I. Nikiforov, 1990, "A New Approach to Deriving Community Exposure Guidelines from No-Observed-Adverse-Effect-Levels," *Regulatory Toxicology and Pharmacology*, 11:314–330.
- Linder, R. E., T. B. Baines, and R. D. Kimbrough, 1974, "The Effect of Polychlorinated Biphenyls on Rat Reproduction," *Food and Cosmetic Toxicology*, 12:63–77.
- Looney, B. B., Grant, M. W., and King, C. M., 1987, Estimation of Geochemical Parameters for Assessing Subsurface Transport at the Savannah River Plant, Savannah River Laboratory Environmental Information Document DPST-85-904, March.
- Ludwig, D. F., Frantzen, K., Friello, P., Kester, J., and Banton, M. I., 1993, An Approach to Toxicity Reference Values for Ecological Risk Assessment. Presented at the 14th Annual Meeting, Society of Environmental Toxicology and chemistry, 14-18 November 1993, Houston, TX.
- Manzo, L., L., et al., 1992, "Metabolic Studies as a Basis for the Interpretation of Metal Toxicity," Toxicology Letters, 64/65:677-686.
- Marie, P. J., et al., 1986, "Effect of Low Doses of Stable Sr on Bone Metabolism in Rats," *Miner. Electroyte Metab.*, 11:5-13.
- Markham, O. D. et al., 1982, "Radionuclides in Pronghorn Resulting from Nuclear Fuel Reprocessing and Worldwide Fallout," *Journal of Wildlife Management*, 46, pp. 30–42.
- Markham, O. D., and D. K. Halford, 1982, "Radionuclides in Mourning Doves Near a Nuclear Facility Complex in Southeastern Idaho," *Wilson Bulletin*, 94, pp. 185–197.
- Markham, O. D., D. K. Halford, and R. E. Autenrieth, 1980, "Strontium-90 Concentrations in Pronghorn Antelope Bones Near a Nuclear Fuel Reprocessing Plant," *Health Physics*, 38:811–816.
- Markham, O. D., R. L. Dickson, and R. E. Autenrieth, 1979, "Plutonium in the Lungs of Pronghorn Antelope Near a Nuclear Fuel Reprocessing Plant," *Health Physics*, 37, pp. 398–400.
- Maughn, J. T., 1993, Ecological Assessment of Hazardous Waste Sites, Van Nostrand Reinhold, New York, NY, 352 pp.

- McConnell, E. E., 1985, "Comparative Toxicity of PCBs and Related Compounds in Various Species of Animals," *Environmental Health Perspectives*, 60:29–33.
- McCormick, S. H., et al., 1997, Work Plan for Waste Area Group 4, Operable Unit 4-13 Comprehensive Remedial Investigation/Feasibility Study, DOE/ID-10550, March.
- McFarland, V. A. and J. U. Clarke, 1989, "Environmental Occurrence, Abundance, and Potential Toxicity of Polychlorinated Biphenyl Congeners: Considerations for a Congener-specific Analysis" *Environ, Health Perspect.*, 81:225–239.
- Montgomery, J. H. and L. M. Welkom, 1990, *Groundwater Chemicals Desk Reference*, Lewis Publishers, Chelsea, MI.
- Morris, R. C., 1998, Potential Use of Habiats Within and Surrounding Facilties at the Idaho National Engineering and Environmental Laboratory by Sensitive Species: A Biological Assessment, Draft ESRF-026, Environmental Science and Research Foundation, Idaho Falls, ID,
- Nagy, K. A., 1987, "Field Metabolic Rate and Food Requirement Scaling in Mammals and Birds," *Ecological Mono.*, 57:111–128.
- National Academy of Science (NAS), 1980, Mineral Tolerance of Domestic Animals, Washington, D.C.
- National Academy of Sciences (NAS), 1977, Arsenic. Washington, D.C.
- National Cancer Institute (NCI), 1977, Bioassay of 1,1,1,-trichloroethane for Possible Carcinogenicity, National Cancer Institute Technical Report Series 3, NTIS PB265082.
- National Cancer Institute (NCI), 1979, "Bioassay of P-chloroaniline for Possible Carcinogenicity," NCI Carcinogenesis Tech. Rep. Ser. No. 189, NTIS PB 295896.
- National Research Council of Canada, (NRCC), 1978, Effects of Arsenic in the Canadian Environment, National Research Council Canada Publication Number NRCC 15391, 349 pp.
- National Toxicology Program (NTP), 1986, NTP Technical Report on the Toxicology and Carcinogenesis of Xylene (Mixed) (60.2% m-xylene, 13.6% p-Xylene, 17.0% Ethylbenzene, and 9.1% o-Xylene) (CAS No. 1330-20-7) in F344N/N Rats and B6C3F1 Mice (Gavage Studies), NIH Publication No.86-2583, Research Triangle Park, N.C.
- National Toxicology Program (NTP), 1989, "Toxicology and Carcinogenesis Studies of Toluene in F/344 Rats and B6C3F1 Mice," *Technical Report Series No. 371*, Research Triangle Park, NC.
- Nekipelov, M. K., 1966, "Hygienic Standard for Aluminum Nitrate in Water Basins." *Hyg. Sanit.* (USSR), 31:204, as cited in National Academy of Sciences, 1980. Mineral Tolerance of Domestic Animals. Washington, DC.
- O'Flaherty, E. J., 1993, "A Pharmacokinetic Model for Chromium," Toxicology Letters, 68:145-158.
- Olson, G. L., D. J. Jeppesen, and R. D. Lee, 1995, *The Status of Soil Mapping for the Idaho National Engineering Laboratory*, INEL-95/0051, Idaho Falls, ID, January.

- Opresko, D. M., B. E. Sample, and G. W. Suter, II, 1995, *Toxicological Benchmarks for Wildlife: 1995 Revision*, ES/ER/TM-86/R2, Oak Ridge National Laboratory, Oak Ridge, TN.
- Ort, J. F. and J. D. Latshaw, 1978, "The Toxic Level of Sodium Selenite in the Diet of Laying Chickens," *Journal of Nutrition*, 108:1114–1120.
- Osborn, D. W. J. Eney, and K. R. Bull, 1983, "The Toxicity of Trialkyl Lead Compounds to Birds," Environmental Pollution, 31A:261-275. As cited in Eisler, 1988.
- Outridge, P. M. and A. M. Scheuhammer, 1993, "Bioaccumulation and Toxicology of Chromium: Implications for Wildlife," *Review of Environmental Contamination and Toxicology*, 130:31–77.
- Penrose, W. R., 1974, "Arsenic in the Marine and Aquatic Environments: Analysis, Occurrence and Significance," CRC Critical Review of Environmental Control, 4:465.
- Perry, H. M. et al., 1983, "Cardiovascular Effects of Chronic Barium Ingestion," *Proceedings of the 17th Annual Conference on Trace Substances in Environmental Health*, Vol. 17, University of Missouri Press, Columbia, MO.
- Perry, H. M. et al., 1985, "Barium-induced Hypertension," Adv. Mod. Environ. Toxicol. Inorga Drinking Water Cardio Vaic Dis, 9:221-229.
- Perry, H. M. Jr. et al., 1989, "Hypertension and Associated Cardiovascular Abnormalities Induced by Chronic Barium Feeding," *Journal of Toxicology and Environmental Health*, 18(3):373–388.
- Pritzl, M. C. et al., 1974, "The Effect of Dietary Cadmium on Development of Young Chickens," *Poultry Science*, 53:2026–2029.
- Puzanova, L., M. Doskocil, and A. Doubkova, 1978, "Distribution of the Development of Chick Embryos after the Administration of Beryllium Chloride at Early Stages of Embryogenesis," *Folia Morphologica*, 27(3):229–231.
- Rai, O, L. E. Eary and J. M. Zachara, 1989, "Environmental Chemistry of Chromium," *The Science of Total Environment*, 86:15–23.
- Reynolds, T. D. et al., 1986, "Vertebrate Fauna of the Idaho National Engineering Laboratory," *Great Basin Naturalist*, 46:513–527.
- Rood, S. M., G. A. Harris, and G. J. White, 1995, Background Dose Equivalent Rates and Surficial Soil Metal and Radionuclide Concentrations for Idaho National Engineering Laboratory, INEL-94/0250, Rev. 0, February
- Rosenfeld, I. and O. A. Beath, 1954, "Effect of Selenium on Reproduction in Rats," *Procedures of the Society of Experimental Biological Medicine*, 87: 295–297.
- Rosomer, G. L. et al., 1961, "Toxicity of Cadmium and Chromium for the Growing Chick," *Poultry Science*, 40:1171–1173.
- RTI (Research Triangle Institute), 1987, Two Generation Reproduction and Fertility Study of Nickel Chloride Administered to CD Rats in Drinking Water: Fertility and Reproductive Performance of

- the Po Generation (Part II of III) and F1 Generation (Part III of III), Final study report. Report submitted to Office of Solid Waste Management, U.S. EPA, Washington, DC.
- Rungby, J. and G. Danscher, 1983a, Localization of Exogenous Silver in Brain and Spinal Cord of Silver Exposed Rats, *Acta Neuropathologica*, 60(1-2):92–98.
- Rungby, J. and G. Danscher, 1983b, "Neuronal Accumulation of Silver in Brains of Progeny from Argyric Rats," *Acta Neuropathologica*, 61(3-4):258-262.
- Rungby, J. and G. Danscher, 1984, "Hypoactivity in Silver-exposed Mice," *Acta Pharmacol and Toxicol*, 55(5):398–401.
- Rungby, J. et al., 1987, "A Quantitative Evaluation of the Neurotoxic Effect of Silver on the Volumes of the Components of the Developing Rat Hippocampus," *Toxicology*, 43(3):261-8.
- Rungby, J., 1990, "An Experimental Study on Silver in the Nervous System and on Aspects of its General Cellular Toxicity," *Dan. Medical Bulletin*, 37(5):442–449.
- Safe, S., 1992, Toxicology, Structure-function Relationships, and Human and Environmental Health Impacts of Polychlorinated Biphenyls: Progress and Problems, *Environmental Health Perspective*, 100:259–268.
- Sattelmacher, P. G., 1962, *Methamoglobinamie durch Nitrate im Trinkwasser*, Schriftenreie Verein Wasser Boden Lufthyg, Berlin-Dahlem No. 20, Gustav Fischer Verlag, Stuttgart (German).
- Sax, N. I. and R. J. Lewis, Jr, 1987, *Hawley's Condensed Chemical Dictionary*, Eleventh Edition. Van Nostrand Reinhold Company, New York, NY, 1288 pp.
- Scheuhammer, A. M., 1987, "The Chronic Toxicity of Aluminum, Cadmium, Mércury, and Lead in Birds: A Review," *Environmental Pollution*, 46(4):263B296.
- Schmalz, B. L., 1972, Radionuclide Distribution in Soil Mantle of the Lithosphere as a Consequence of Waste Disposal at the National Reactor Testing Station, IDO-10049.
- State of Idaho, 1996, Guidance Document for Petroleum Releases, Division of Environmental Quality Remedial Bureau, Idaho DEQ-RBCA Document No. 230.
- Schroeder and Mitchener, 1975, "Life-term Effects of Mercury, Methylmercury and Nine Other Trace Metals on Mice," *Journal of Nutrition* 105:452.
- Schroeder, H. A. and J. J. Balassa, 1967, "Arsenic, germanium, tin and vanadium in mice: Effects on growth, survival and tissue levels," Journal of Nutrition, 92:245–252.
- Schroeder, H. A. et al., 1968, Zirconium, Niobium. Antimony, Vanadium, and Lead in Rats: Life Term Studies, J. Nutr. 95: 95-1101.
- Schultz, V. and F. W. Whicker, 1982, Radioecology: Nuclear Energy and the Environment, CRC Press, Inc., Boca Raton, FL.

- Scow, K., et al., 1981, Exposure and Risk Assessment for Silver, Report to U. S. Environmental Protection Agency, Office of Water Regulations and Standards, Washington, D.C., by Arthur D. Little, Inc., Cambridge, MA, PB85-211993.
- Shacklette H. T. And J. C. Boerngen, 1984, Element Concentrations in Soils and Other Surficial Materials of the Conterminous United States, USGS Profession Paper 1270.
- Sheppard, M. I. and Thibault, D. H., 1990, "Default Soil Solid/Liquid Partition Coefficients, Kds, for Four Major Soil Types: A Compendium," *Health Physics*, 59(4):471–482.
- Shinogi, M. and S. Maeizumi, 1993, "Effect of Preinduction of Metallothioncin on Tissue Distribution of Silver and Hepatic Lipid Peroxidation," *Biological and Pharmacological Bulletin* 16(4): 372–374.
- Shleien, B., 1992, *The Health Physics and Radiological Health Handbook*, Scinta Publishers, Silver Springs, MD.
- Shopp et al., 1984, "Naphthalene Toxicity in CD-1 Mice: General toxicology and Immunotoxicology," Fundamentals of Applied Toxicology, 4:406–419.
- Simon, C. H. Manzke, H. Kay, and G. Mrowetz, 1964, Occurrence, Pathogenesis, and Possible Prophylaxis of Nitrite Induced Methemoglobinemia, Zeitschr. Kinderheilk, 91:124–138 (German).
- Sittig, M., 1981, Handbook of Toxic and Hazardous Chemicals, Noyes Publications, Park Ridge, NJ.
- Skoryna, S. C., and M. Fuskova, 1981, "Effects of Stable Strontium Supplementation," *Handbook of Stable Strontium*, S. C. Skoryna, ed., Plenum Press, p. 593–613.
- Sleight, S. D. and O. A. Atallah, 1968, "Reproduction in the Guinea Pig as Affected by Chronic Administration of Potassium Nitrate and Potassium Nitrite," *Toxicol. Appl. Pharmacol.*, 12:179-185.
- Smith, G. J. et al., 1988, "Reproduction in Black-crowned Night Herons Fed Selenium," Lake and Reservoir Management. 4(2):175–180.
- Smith, I. C. and B. L. Carson, ed., 1981. Trace metals in the environment. Volume 6: Cobalt an appraisal of environmental exposure. Ann Arbor, MI: Ann Arbor Science Publishers, Inc.
- Smyth, H. F., Jr. and C. P. Carpenter, 1948, "AFurther Experience with the Range Finding Test in the Industrial Toxicology Laboratory," *Journal of Industrial Hygiene and Toxicology*, 30:63–68.
- Steele, M. J. et al., 1990, "Assessing the Contribution from Lead in Mining Wastes to Blood Lead," *Regulatory Toxicology and Pharmacology*, 11:158–190.
- Steven, J. D. et al., 1976, Effects of Chromium in the Canadian Environment, NRCC No. 15017, National Resources Council, Ottawa, Canada.
- Storey, E., 1961, "Strontium Rickets, Bone Calcium, and Strontium Changes," Austral. Ann. Med., 10:213–222.

- Suter, G. W. and C. L. Tsao, 1996, Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota: 1996 Revision, Risk Assessment Program, Oak Ridge National Laboratory, June, ES/ER/TM-96/R2.
- Suter, G. W. II, 1993, Ecological Risk Assessment, Lewis Publishers, Chelsea, Michigan, 538 pp.
- Suter, G. W., II, 1989, Chapter 2, "Ecological Endpoints," *Ecological Assessments of Hazardous Waste Sites: A Field and Laboratory Reference Document*, W. Warren-Hicks, B. R. Parkhurst, and S. S. Baker (eds.), U.S. Environmental Protection Agency, March, EPA 1600/3-89/013.
- Suter, G. W., II, M. A. Futrell, and G. A. Kerchner, 1992, Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Aquatic Biota on the Oak Ridge Reservation, Oak Ridge, Tennessee, Energy Systems Environmental Restoration Program, ORNL Environmental Restoration Program, Oak Ridge National Laboratory, September, ORNL/ER-139 ES/ER/TM-96.
- Suter, G. W., II, M. E. Will, and C. Evans, 1993, Toxicological Benchmarks for Screening Potential Contaminants of Concern for Effects on Terrestrial Plants, Environmental Restoration Division, ORNL Environmental Restoration Program, Oak Ridge National Laboratory, September, ES/ER/TM-85.
- Teratology, 1988, 38:253.
- Thompson, A, S. L. Hansard, and M. C. Bell, 1959, "The Influence of Aluminum and Zinc Upon the Absorption and Retention of Calcium and Phosphorus in Lambs," *Journal of Animal Science*, 18:187, as cited in National Academy of Sciences, 1980, *Mineral Tolerance of Domestic Animals*, Washington, D.C.
- Thurston, H., G. R. Gilmore, and J. E. Swales, 1972. "Aluminum Retention and Toxicity in Renal Failure," *The Lancet*, 1: 881-883.
- Tooele Army Depot: North Area Draft Sitewide Ecological Risk Assessment, Volume 1, SFIM-AEC-IR-CR-9, January 1996.
- Torkelson, T. R. et al., 1958, "Toxicity of 1,1,1-trichloroethane as Determined on Laboratory Animal and Human Subjects," Am. Ind. Hyg. Assoc. J, 19:353-362.
- TOXNET, 1994, National Library of Medicine On-line Computer Database.
- Travis, C. C. and A. D. Arms, 1988, "Bioconcentration of Organics in Beef, Milk, and Vegetation," Environmental Science and Technology, 22:271–274.
- Trivedi, B. et al., 1989, "Embryotoxicity and Fetotoxicity of Orally Administered Hexavalent Chromium in Mice," *Reproductive Toxicology*, 3(4):275–278.
- Turk, J. L. and F. H. Kratzer. 1960. The effects of cobalt in the diet of chicks. Poult. Sci. 39:1302. (Abstract).
- Ungvary, G. et al., 1980. "Studies on the Embryotoxic Effect of Ortho-, Meta-, and Para-xylene," *Toxicology*. 18:61–74.

- Uthus, E. O., 1992, "Evidence for Arsenic Essentiality," *Environmental Geochemistry and Health*, 14(2):55-58.
- Valdivia, R. et al., 1978, Effect of Dietary Aluminum on Animal Performance and Tissue Mineral Levels in Growing Steers, *Journal of Animal Science*, 47:1351, as cited in National Academy of Sciences, 1980, Mineral Tolerance of Domestic Animals. Washington, D.C.
- Van Bruwaene, R. et al., 1981, "Metabolism of Antimony-124 in Lactating Dairy Cows," Health Physics.
- Van Zinderen Bakker and J. F. Jaworski. 1980. Effects of vanadium in the Canadian environment. Ottawa, Canada: National Research Council Canada, Associate Committee Scientific Criteria for Environmental Quality. 1–94.
- VanHorn, R., N. L. Hampton, and R. C. Morris, 1995, Guidance Manual for Conducting Screening Level Ecological Risk Assessment at the INEL, Idaho National Engineering Laboratory, EG&G Idaho, Inc., Idaho Falls, ID, INEL-95/0190, April.
- Venugopal, B. and T. D. Luckey, 1978, Metal Toxicity in Mammals, Plenum Press, New York.
- Verschueren, 1983, Environ. Data Org. Chem., p. 619.
- Walker, E. D., 1994, Stormwater Management at the Arid INEL, EGG-M-94112.
- Wallace, A., G. V. Alexander, and F. M. Chaudhry, 1977, "Phytotoxicity of Cobalt, Vanadium, Titanium, Silver and Chromium," *Commun Soil Sci Plant Anal*, 8(9):751–756.
- Walton, G., 1951, Survey of Literature Relating to Infant Methemoglobinemia Due to Nitrate-Contaminated Water, Am. J. Public Health, 41:986–996.
- Weber, C. W. and B. L. Reid, 1968, "Nickel Toxicity in Growing Chicks," *Journal of Nutrition* 95:612-616.
- Wehrli, B. and W. Stumm. 1989. Vanadyk in natural waters: Adsorption and hydrolysis promote oxygenation. Geochim Cosmochim Acta 53:69–77.
- Weir, R. J. Jr. and R. S. Fisher, 1972, "Toxicologic Studies on Borax and Boric Acid," *Toxicol. Appl. Pharmacol.*, 23:351–364.
- Weitz, A. and K. E. Ober, 1965, "Physiological Distribution of Antimony after Administration of Sb-124 Labeled Tartar Emetic," *Bulletin of the World Health Organization*, 33:137–142.
- White, D. H. and M. P. Dieter, 1978, "Effects on dietary vanadium in mallard ducks," Journal of Toxicology and Environmental Health.
- Wilber, C. G., 1980, "Toxicology of Selenium: A Review," Clinical Toxicology 17:171–230.
- Will, M. E., Suter II, G. W., 1995, Toxicology Benchmarks for Screening Potential Contaminants of Concern for Effects on Terrestrial Plants, September.

- Wills, J. H., G. E. Groblewski, F. Coulston, 1981. "Chronic and Multigeneration Toxicities of Small Concentrations of Cadmium in the Diet of Rats." *Ecotoxicology and Environmental Safety*, 5:452-464.
- WINCO, 1993, Track 1 Decision Documentation Package Waste Area Group 3 Operable Unit 02 Site CPP-66 CPP CFSGF Ash Pit, Idaho National Engineering Laboratory, Idaho Falls, ID, September.
- WINCO, 1994a, Track 1 Decision Documentation Package Waste Area Group 3 Operable Unit 02 Site CPP-65 CPP Sewage Treatment Plant, Idaho National Engineering Laboratory, Idaho Falls, ID, April.
- WINCO, 1994b, Track 1 Decision Documentation Package Waste Area Group 3 Operable Unit 03 Site CPP-67 CPP Percolation Ponds #1 and #2, Rev. 3, Idaho National Engineering Laboratory, Idaho Falls, ID, April.
- Witmer, C. M., R. Harris, and S. I. Shupack, 1991, "Oral Bioavailability of Chromium From a Specific Site," *Environmental Health Perspective*, 92:105–110.
- Wobeser, G. A., N. O. Nielson, and B. Schiefer, 1976, "Mercury and mink. 2. Experimental Methylmercury Intake in Mink," *Canadian Journal of Comparative Medicine*, 40:34–45.
- World Health Organization (WHO), 1988, "Vanadium." Environmental Health Criteria 81. World Health Organization, Geneva, Switzerland.
- WHO (World Health Organization), 1983, "Selected Radionuclides," Environmental Health Criteria, 25, Geneva, Switzerland.
- Wren, C. D. et al., 1987, "The Effects of Polychlorinated Biphenyls and Methylmercury, Singly and in Combination, on Mink, II: Reproduction and Kit Development," *Archives of Environmental Contamination and Toxicology*, 16:449-454.
- Wren, C. D., 1986, "A Review of Metal Accumulation and Toxicity in Wild Mammals, I. Mercury," Environmental Research, 40(1):210B44.
- Wynder, E. L. and D. Hoffman, 1959, "The Carcinogenicity of Benzofluoranthenes," *Cancer*, 12:1194-1199.
- Zmudzki, J. et al., 1983, "Lead Poisoning in Cattle: Reassessment of the Minimum Toxic Oral Dose," Bulletin of Environmental Contaminants, 30:435-441.